



King's Research Portal

DOI:

[10.1016/j.jaut.2018.02.009](https://doi.org/10.1016/j.jaut.2018.02.009)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Lewis, M. J., McAndrew, M. B., Wheeler, C., Workman, N., Agashe, P., Koopmann, J., Uddin, E., Morris, D. L., Zou, L., Stark, R., Anson, J., Cope, A. P., & Vyse, T. J. (2018). Autoantibodies targeting TLR and SMAD pathways define new subgroups in systemic lupus erythematosus. *Journal of Autoimmunity*. <https://doi.org/10.1016/j.jaut.2018.02.009>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Figure S1. Flow chart of patients




Figure S2. Principal component analysis of discovery and validation datasets




Figure S3. Reduced subspace heatmap showing differences between in autoantibody levels across four SLE subgroups in discovery and validation datasets




Figure S4. Principal component analysis of autoantibody levels in SLE individuals and controls




Figure S4





Figure S5. Distribution of autoantibody positivity by ancestry

Stacked barchart showing distribution of ancestry across SLE patients for each autoantibody. P value calculated by Fisher exact test, with FDR correction.




Figure S6. Distinctive SLE autoantibodies profiles according to ancestry

(A) Differential antibody positivity according to ancestry, showing autoantibodies with increased positivity in Afro-Caribbean (AA) SLE, Hispanic (HA) SLE and European (EA) ancestry SLE. Autoantibodies were selected as statistically different between SLE ancestral groups by one-way ANOVA (FDR corrected $P < 0.01$). (B) Boxplots of key autoantigens specific for different ancestries in SLE. (C) Linear discriminant analysis based on plots showing separation of Afro-Caribbean, Hispanic and European ancestry SLE individuals according to linear discriminant factors LD1, LD2 and LD3. Right hand panel, 3-dimensional plot of LD1, LD2 and LD3. (D) Table of LDA predicted categorisation of SLE ancestral groups and control individuals compared to observed groupings.




Figure S7. Effect of HLA alleles on autoantibody levels in SLE

Autoantigens colour-coded by cluster (red=1a, purple=1b, green=2, blue=3). P values (without adjustment for multiple testing) were calculated by fitting a linear regression model including antibody titre and genetic population structure PC1-4. Dark blue dots represent antibody positivity defined as >2 SD of control population.




Figure S8. Cross-validation of elastic net binomial regression model

10-fold cross-validation was performed on the discovery dataset to tune L1-L2 penalisation parameter α and shrinkage parameter λ by minimising binomial deviance.





Figure S9. SLE cluster heatmap of binomial regression model autoantibodies




Figure S10. ROC curve analysis of mixture discriminant analysis model incorporating ANA and dsDNA antibodies




Figure S11. Canonical variate plot of mixture discriminant analysis model

Canonical variate plots from 26-autoantibody mixture discriminant analysis (MDA) model showing 4 SLE clusters and 1 control cluster in (A) discovery and (B) validation datasets. Cluster centroids are marked C for 1 control centroid and S for 4 SLE centroids.




Figure S12. Heatmap of autoantibody levels grouped by ancestry for 26-autoantibody multinomial regression model

